

<b>T1</b>	<ol style="list-style-type: none"><li>1. <math>d_0 =</math></li><li>2. <math>d_{ISM} =</math></li></ol>
<b>T2</b>	<ul style="list-style-type: none"><li>• <math>E_{tot} =</math></li><li>• <math>\frac{v_p}{v_a} =</math></li></ul>
<b>T3</b>	<ol style="list-style-type: none"><li>1. <math>\lambda_{Kathmandu} =</math></li><li>2. <math>t_{Kathmandu} =</math></li><li>3. <ol style="list-style-type: none"><li>i. <math>\Delta t =</math></li><li>ii. <math>t_{Ranchi} =</math></li></ol></li></ol>
<b>T4</b>	<ol style="list-style-type: none"><li>1. <math>M =</math></li><li>2. <math>d_{var} =</math></li></ol>

<b>T5</b>	$a_{sat} =$
-----------	-------------

<b>T6</b>	<ol style="list-style-type: none"><li>1.</li><li>2.</li></ol>
-----------	---

<b>T7</b>	<ol style="list-style-type: none"><li>1. <math>\lambda_{obs} =</math></li><li>2. <math>m_{corr} =</math></li><li>3. <ol style="list-style-type: none"><li>i. <math>F_{app} =</math></li><li>ii. <math>F_{corr} =</math></li></ol></li><li>4. <math>S_{app} =</math></li><li>5. <math>r_{gal} =</math></li></ol>
-----------	---

<b>T8</b>	<ol style="list-style-type: none"><li>1. <math>t_{rise} =</math></li><li>2. <math>\Delta\theta =</math></li></ol>
-----------	---

<b>T9</b>	<p>1. <math>t_{return} =</math></p> <p>2. <math>v_{esc} =</math></p>
-----------	--

<b>T10</b>	<p>1. i.</p> <p>ii.</p> <p>2. i. <math>m_1 = M_{\odot}</math></p> <p>ii. <math>m_2 = M_{\odot}</math></p> <p>iii. Upper Limit / Lower Limit / Exact</p>
------------	---

<b>T11</b>	<p>1. <math>\rho_c =</math></p> <p>2. <math>\rho_c =</math></p> <p>3. <math>N =</math></p>
------------	--

<p><b>T12</b></p>	<p>1. <math>\lambda_{max} =</math></p> <p>2. <math>r_{cloud} =</math></p> <p>3. <math>\theta =</math></p> <p>4. <math>D_{tele} =</math></p>
-------------------	---

## T1

1. (a) (1 point) We know  $m_t - m_e = -2.5 \log \left( \frac{F_t}{F_e} \right)$

(1 point) since,  $F_{lim} \propto \frac{1}{D^2}$  so  $m_t - m_e = 5 \log (1000/7)$

$$m_t = 10.8 + 6$$

(1 point)  $m_t = 16.8$

Now we know  $m = M + 5 \log \left( \frac{r}{10pc} \right)$

$$\frac{r}{10} = 10^{3.454}$$

(2 points) So  $r = 29Kpc$  is the maximum distance till which student can detect the star using telescope.

- (b) (2 points)  $m' = m + A$ , thus  $m' = 16.8 + \alpha r$

(1 point) Now,  $m' = M + 5 \log \left( \frac{r}{10pc} \right)$   $16.8 + 0.00005 * r = -0.5 + 5 \log(r) - 5$

(2 points)  $r = 10^{4.46 - 0.00001r}$

This equation can't be solve analytically.

(2 points) So, if we do the iteration for this, taking initial value from previous result of 29Kpc, which is good approximation.

(3 points) Then, we get  $r = 18.73Kpc$ .

## T2

2. (2 points)  $r_p = a(1 - e)$  is the perihelion distance.  $r_a = a(1 + e)$  is the aphelion distance. Use the law of conservation of energy to get:

(2 points)

$$E = -\frac{GMm}{r_p} + \frac{mv_p^2}{2} = -\frac{GMm}{r_a} + \frac{mv_a^2}{2}$$

(1 point) From the conservation of angular momentum,

$$mv_a r_a = mv_p r_p$$

$$\implies v_a = v_p \frac{r_p}{r_a}$$

(3 points) Substitute velocities and rearrange the first equation to get (alternative expression making  $v_p$  the subject is also possible and correct):

$$v_a^2 \left( 1 - \frac{r_a^2}{r_p^2} \right) = 2GM \left( \frac{1}{r_a} - \frac{1}{r_p} \right)$$

(3 points) Then, substitute this into the equation for the total mechanical energy:

$$E = \frac{-GMm}{r_a} + \frac{GMmr_p^2}{r_p^2 - r_a^2} \left( \frac{r_p - r_a}{r_a r_p} \right)$$

Which gives (after simplifying or any equivalent formulation):

$$E = \frac{-GMm}{2a} \left( \frac{2}{1+e} - \frac{1-e}{1+e} \right)$$

(2 points) Finally giving us:

$$E = -\frac{GMm}{2a}$$

(2 points) Substitute  $r_a$  and  $r_p$  in the energy equations to get:

$$\frac{v_a}{v_p} = \frac{1-e}{1+e}$$

(or use the conservation of angular momentum directly).

### T3

3. (a) (3 points)

$$lon_g = \frac{5.75}{12} \times 180 = 86.25 \text{ deg } E$$

(4 points)

$$lon_k = lon_g - \frac{105 \times \cos \phi}{6378} \times \frac{180}{\pi} = 85.415 \text{ deg } E$$

(b) (3 points)

$$t_{diff} = \frac{105 \times \cos \phi}{6378} \times \frac{12 \times 3600}{\pi} \text{ s} = 200.4 \text{ s} = 3 \text{ min } 20.4\text{s}$$

$$t_{ktm} = 8:35 \text{ pm} - t_{diff} = 8:31:40$$

(1 point) Answer can be written as 8:32pm, or 8:31:40pm.

- (c) (2 points) Ravi has the same local time as Kathmandu. On his watch, however, his time zone is UTC +5:30. Thus, the difference will be  $15 - t_{diff}$  mins, which is 11 mins 40 (39.6) seconds.

(2 points) As per his watch, therefore, the occultation will occur at 8:16:40 pm, or 8:17 pm.

## T4

4. (a) (2 points)  $L \propto P^{2.5} \rightarrow L = Kp^{2.5}$ , taking log on both side then  $\log L = \log K + 2.5 \log P$

$$\log L = \log F + \log 4 + \log \pi + 2 \log r$$

(1 points)  $m_1 - m_2 = -2.5 \log \left( \frac{F_1}{F_2} \right)$

For ease of calculation we can use Vega as reference star and  $m$  be apparent magnitude of cepheid variable then

(1 point)  $\frac{-m}{2.5} = \log F_1 - \log F_o$

$$m = M + 5 \log \left( \frac{r}{10 \text{pc}} \right)$$

Solving these equation we get,

(3 points)

$$M = -6.25 \text{Log} P + C$$

,where  $C = -2.5(\text{Log} K - \text{Log} 4 - \text{Log} F_o - \text{Log} \pi) + 5 \text{Log} 10$

- (b) (1 point) So for above equation of magnitude luminosity relation  $C = -1.5$ ,

(1 point) we suppose,  $m$  is the average magnitude for luminosity,  $\frac{1}{2}(L_{\max} + L_{\min})$ .

(1 point) Now  $m - m_{ref} = -2.5 \log \left( \frac{\frac{1}{2}(L_1 + L_2)}{L_{ref}} \right)$

$$m - m_{ref} = 2.5 \log 2 - 2.5 \log \left( \frac{L_1 + L_2}{L_{ref}} \right)$$

$$m = m_{ref} + 2.5 \log 2 - 2.5 \log \left( \frac{L_1 + L_2}{L_{ref}} \right)$$

(1 point)  $\frac{L_{tot}}{L_{ref}} = \left( \frac{L_1 + L_2}{L_{ref}} \right) = \left( \frac{L_1}{L_{ref}} + \frac{L_2}{L_{ref}} \right)$   
 $= (10^{-0.4m_1} + 10^{-0.4m_2}) 10^{0.4m_{ref}}$

(1 point) Now substituting the values,  
 $m = 2.5 \cdot \log 2 - 2.5 \cdot \log(10^{-\frac{2}{5}m_1} + 10^{-\frac{2}{5}m_2})$

(1 point) From graph  $m_{max} = 28$  and  $m_{min} = 22$  thus  $m = 22.75$  and period is 20 days.

(1 point) So, using the Absolute magnitude relation with period we obtained above we can write  $M = -9.63$ .

(1 point) Substitute the value in this equation  $m = M + 5 \log \left( \frac{r}{10pc} \right)$  we get,  $r = 29.92$  Mpc.

## T5

(3 points) Firstly, calculate the geocentric orbit height by using Newton's law of gravitation and setting  $T = 23$  hours 56 mins 4 secs.

$$\frac{GMm}{d^2} = \frac{mv^2}{d} \text{ and } v = \frac{2\pi d}{T}$$

(1 point) Solve for  $d$  to get:

$$d = \left( \frac{GMT^2}{4\pi^2} \right)^{1/3}$$

(1 point) Substitute the correct values to obtain  $d = 4.216 \times 10^7$  m.

(5 points) Then, construct a triangle, such that from the law of sines we have:

$$\frac{\sin(90 + a)}{d} = \frac{\sin(90 - (\phi + a))}{r}$$

where  $r$  is the radius of the Earth,  $a$  is the altitude and  $\phi$  is the latitude of Kathmandu.

(2 points) Simplifying:

$$\cos(a) \frac{r}{d} = \cos(\phi) \cos(a) - \sin(\phi) \sin(a) \implies 1 - \tan(\phi) \tan(a) = \frac{r}{d \cos(\phi)}$$

(2 points) Rearranging and solving for  $a$ , you get:

$$a = \tan^{-1} \left( \frac{1 - \frac{r}{d \cos(\phi)}}{\tan(\phi)} \right)$$

(1 point) Plugging in the values:

$$a = 57.64^\circ$$

## T6

1. (1 point) The mean synodic period of the Moon is 29.53059 days. Dashain happened 12 synodic periods earlier, so  
 (3 points)  $12 \times 29.53059 = 354.36708$  days = 354 days 8 hours 48 minutes.  
 (1 point) Because 2024 is a leap year, there are 366 days between the two 12 Octobers.  
 (3 points) Dashain in 2023 will be 354 days before the 12th of October.  
 (2 point) In other words, it was 12 days after the 12 October, 2023, which was 24 October, 2023.
2. (1 point) Dashain in 2025 will be 354 days 8 hours 48 minutes after 11am on 12 October 2024,  
 (2 points) which is 11 days before 12 October 2025 at around 7:48pm on 1 October 2025.  
 (2 points) 4 days and 12 hours after this is 7:48am on 6 October 2025. Therefore, *Kojagrat Purnima* will fall on 6 October 2025.

## T7

1. (1 point)
 
$$\lambda_{obs} = (1 + z) \times \lambda_{rest} = 3.379 \times 656.3\text{nm}$$
 (1 point)
 
$$= 2217.6\text{nm}$$
2. (2 points) Lensing is multiplicative on the flux. So:

$$m_{corr} = m_{obs} + 2.5 \log(30) = 24.19$$

3. (2 points)

$$f_{obs} = 3631 \times 10^{-\frac{20.5}{2.5}} \text{Jy} = 2.29 \times 10^{-5} \text{Jy}$$

- (2 points)

$$f_{corr} = \frac{f_{obs}}{\mu} = 7.64 \times 10^{-7} \text{Jy}$$

4. (2 points)  $\Sigma_{UV} = f_{obs}/13$ .

(2 points) In magnitudes,  $\Sigma'_{UV} = -2.5 \log(\Sigma_{UV}/3631) = 23.3 \text{ mag arcsec}^{-2}$

5. (3 points)

$$r = \sqrt{13 \times 8^2 / \pi \mu} = 3 \text{ kpc}$$

[1 point for correct placement of magnification factor, 1 for squaring scale, 1 for correct answer.]

## T8

1. (3 points) We know:

$$\sin(a) = \sin(\phi) \sin(\delta) + \cos(\phi) \cos(\delta) \cos(H)$$

- (1 point) At rise,  $a = 0$ .

$$\cos(H) = -\tan(\phi) \tan(\delta)$$

(3 points) Solve for H to get  $H = 5 \text{ hours } 49 \text{ minutes } 59 \text{ seconds}$ .

- (2 points)

$$t_{rise} = 12\text{pm} - H = 6:10 \text{ am}$$

2. (1 point) Since the stars are very close, you can use planar geometry to solve this problem.

$$d = \sqrt{(\Delta\alpha)^2 + (\Delta\delta)^2}$$

(2 point) Substitute  $\Delta\alpha = 0.08376 \text{ rad}$

(2 points)  $\Delta\delta = 0.01927 \text{ rad}$

(1 points) Solving in the original equation, obtain  $d = 0.08595 \text{ rad}$  or in degrees,  $4.92 \text{ deg}$ . Answers that round to 4.9 degrees are acceptable. Answers in degrees or subunits there of for any part are acceptable. Correct solutions with spherical trigonometry will also receive full points.

## T9

1. (2 points) Notice that the expression for the acceleration due to gravity becomes:

$$a = -\frac{4}{3} G \pi \rho r$$

which can be expressed as:

$$a = -k^2 r, \text{ where: } k = \sqrt{\frac{4}{3}G\pi\rho}$$

(1 point) This is the equation for a harmonic oscillator, as given in the hint. The time period is given by:

$$T = \frac{2\pi}{k}$$

(2 points) which simplifies to:

$$T = \sqrt{\frac{3\pi}{G\rho}}$$

2. (3 points) From the expression of V, the potential difference between  $r = R$  and  $r = 0$  is given by  $\Delta V_{int} = \frac{GM}{2R}$

(2 points) Finally, to escape:

$$\frac{1}{2}mv_{esc}^2 - \frac{4}{3}Gm\rho\pi R^2 - \Delta V_{int} = 0$$

(5 points) Collecting terms and simplifying in terms of  $v_{esc}$ :

$$v_{esc} = 2R\sqrt{G\pi\rho}$$

## T10

1. (1 point) We know  $V_1 = \omega a_1$  and  $V_2 = \omega a_2$ .

(2 points) From Kepler's Third law:  $P^2 = \frac{4\pi^2}{GM}a^3$ , where  $M = m_1 + m_2$  and  $a = a_1 + a_2$ .

(1 point) We know that  $P = \frac{2\pi}{\omega}$ ,  $a_1 = \frac{m_2}{(m_1+m_2)}a$  and  $a_2 = \frac{m_1}{(m_1+m_2)}a$

(2 points) Replacing above value in Kepler's equation we get,

$$G(m_1 + m_2) = 4\pi^2 \frac{\omega^2}{4\pi^2} \frac{(m_1+m_2)^3}{m_2^3} a_1^3$$

$$\frac{m_2^3}{(m_1+m_2)^2} = (\omega a_1)^3 \frac{P}{2\pi G}$$

(2 points) When we observe the star we can't measure the real orbital velocity but the component of it as the plane of orbit and astronomer line of sight have angle  $\theta$ , thus

(1 point)  $\frac{m_2^3}{(m_1+m_2)^2}(\sin^3\theta) = (v_{1obs})^3\frac{P}{2\pi G}$ , which is required mass function, where  $v_{1obs}$  is maximum observed orbital velocity

(1 point) similarly we can write another mass function,  $\frac{m_1^3}{(m_1+m_2)^2}(\sin^3\theta) = (v_{2obs})^3\frac{P}{2\pi G}$

2. (1 point) We know  $\frac{v_1}{v_2} = \frac{m_2}{m_1}$

(2 points) From graph we can see that  $\frac{v_1}{v_2} = 1.5$  thus  $m_2 = 1.5m_1$ , period is 40 days and  $v_{1obs}$  is 75 km/s, and for minimum value we consider  $\sin\theta$  as 1, if we put these value in mass function, we get

(1 point)  $0.54m_1 = 1.74M_\odot$

$m_1 = 3.2M_\odot$ , similarly

(1 point)  $m_2 = 4.9M_\odot$

## T11

1. (1 point) Use the escape velocity definition:

$$v_{esc} = \sqrt{\frac{2GM}{R}}$$

(1 point) together with Hubble's law:

$$v = H_0R$$

And:

$$M = \frac{4}{3}\pi R^3 \rho_c$$

(4 points) Substitute Hubble's Law and Mass in the escape velocity definition to obtain:

$$\rho_c = \frac{3H_0^2}{8\pi G}$$

2. (2 units)

$$H_0 = 2.2685 \times 10^{-18} \text{ s}^{-1}$$

(2 points) Get:  $\rho_c = 9.20 \times 10^{-27} \text{ kg m}^{-3}$

3. (1 point)

$$M = \frac{4}{3}\pi \rho_c R_\odot^3$$

(1 point)

$$N = \frac{M}{M_{\text{tennis}}}$$

(3 points) Substitute values to get:

$$N = 89.45 \text{ (either 89 or 90)}$$

(both acceptable).

## T12

1. (2 points)

$$\lambda_{\text{max}} T = 2.898 \times 10^{-3} \text{ mK}$$

(1 point) Thus,  $\lambda_{\text{max}} = 57.96 \mu\text{m}$ .

2. (2 points) Use the Stefan-Boltzmann Law and compare to the Sun to get:

$$R_{\text{dust}} = R_{\odot} \sqrt{\frac{L_{\text{dust}}}{L_{\odot}} \frac{T_{\odot}^4}{T_{\text{dust}}^4}}$$

(2 points) Substituting values, you get:

$$R_{\text{dust}} = 120.4 \text{ pc}$$

3. (3 points)

$$d_{\text{dust}} = \frac{cz}{H_0} = \frac{0.01 \times 2.998 \times 10^5}{70} \text{ Mpc} = 42.83 \text{ Mpc}$$

(2 points)


$$\theta = \frac{R_{\text{dust}}}{d_{\text{dust}}} \text{ rad} = 0.580^\circ$$

4. (3 points)

$$D > 1.22 \frac{\lambda'_{\text{max}}}{\theta} = 25.4 \text{ m}$$

$\lambda_{\text{max}}$  needs to have redshift accounted, i.e.  $\lambda'_{\text{max}} = (1+z)\lambda_{\text{max}}$  (or some equivalent). Only 1 point gained if  $\lambda_{\text{max}}$  is used for correct formula.

**OC1 Sky Chart:**



1.	$\phi_{location} =$																						
2.	On the chart.																						
3.	On the chart.																						
4.	<table border="1"> <thead> <tr> <th>Number</th> <th>Constellation Name</th> </tr> </thead> <tbody> <tr><td>C1</td><td></td></tr> <tr><td>C2</td><td></td></tr> <tr><td>C3</td><td></td></tr> <tr><td>C4</td><td></td></tr> <tr><td>C5</td><td></td></tr> <tr><td>C6</td><td></td></tr> <tr><td>C7</td><td></td></tr> <tr><td>C8</td><td></td></tr> <tr><td>C9</td><td></td></tr> <tr><td>C10</td><td></td></tr> </tbody> </table>	Number	Constellation Name	C1		C2		C3		C4		C5		C6		C7		C8		C9		C10	
	Number	Constellation Name																					
	C1																						
	C2																						
	C3																						
	C4																						
	C5																						
	C6																						
	C7																						
	C8																						
	C9																						
C10																							
5.	<table border="1"> <thead> <tr> <th>Code</th> <th>Star</th> </tr> </thead> <tbody> <tr><td>S1</td><td></td></tr> <tr><td>S2</td><td></td></tr> <tr><td>S3</td><td></td></tr> <tr><td>S4</td><td></td></tr> </tbody> </table>	Code	Star	S1		S2		S3		S4													
	Code	Star																					
	S1																						
	S2																						
	S3																						
S4																							
6.	On the chart.																						
7.	$\alpha_{Aldebaran} =$ $\delta_{Aldebaran} =$																						

**OC2**

Asteroids	Dwarf Planets	Satellites

**OC3**


Object	Messier Number (e.g. M32)	Constellation (e.g. ORI or Orion)
O1		
O2		
O3		
O4		
O5		
O6		

## OC1

1. **Answer: 27°7' N**

[ Answer between 27-28° gains all 4 points, with each degree of inaccuracy 0.5 points are lost. e.g. if a student answer 26° or 29°, they would receive 3.5 points]

2. Answer:




[ With 10° of inaccuracy, 0.5 points is lost for cardinal points, the planets have to be circled correctly to gain full points. Naming of the planet is not needed ]

3. Same map as above except for the local meridian to go straightly from North to South.

[ 10 degree of inaccuracy will lead to a 0.5 points loss ]

4. Can be any constellations shown in the sky map below.  
 [ If the tracing composes of at least the brightest stars then the student will earn full points, missing one or more prominent star will lead to a loss of 0.5 point ]




5. Answer:


- A. Alpha Canis Majoris (Sirius)
- B. Alpha Aurigae (Capella)
- C. Beta Orionis (Rigel)
- D. Alpha Canis Minoris (Procyon)

[ 0.5 points for each star can be earned by writing the right Bayer designated name or abbreviation of the star, 0.5 points for each star can be earned if the star is in the right order in the list, 0.5 points for each star can be earned if the star is marked correctly in the sky map ]

**Note:** Sky map is attached below



6. Answer: Sky map is attached. Look for M42, M81, M41, M31,  $\gamma$  Orionis (Ori),  $\alpha$  Persei (Per) and  $\alpha$  Cyg.  
[ If the object labeled is within the size of the circles below, then the student earns full credit. If the object is in the constellation its supposed to be but not in the right place, the student earns 0.5 points ]



7. Answer: Equatorial coordinates of the star Aldebaran ( $\alpha$  Tauri)

**Right Ascension: 4h 36m**

**Declination: 16° 31'**

[ If the RA and Dec is between

3.5h to 5.5h and 15.5° to 17.5°, the student earns full credit (2.5 points).


2.5h to 6.5h and 14.5° to 18.5° = 2 points,

1.5h to 7.5h and 13.5° to 19.5° = 1.5 points,

0.5h to 8.5h and 12.5° to 20.5° = 1.0 points,

-0.5h to 9.5h and 11.5° to 21.5° = 0.5 points,

Anything beyond this range will not earn any credit. ]




**OC2(1.5x6)**

Asteroids	Dwarf Planets	Satellites
5	2	1,3,4,6

**OC3(1.5x12)**

	M 97	Ursa Major
	M101	Ursa Major
	M51	Canes Venatici
	M45	Taurus



M1

Taurus



M42/M43 both will get full marks

Orion

## OT1

1.	
2.	On the Image Above
3.	On the Image Above.

Based on your observations, complete the following table:

Age of the Moon (No. of days since new moon)	
Hemisphere from where the Moon is being watched	
Official IAU name of the constellation in which the Moon is located	

## OT2


1.	<input type="checkbox"/> Mercury <input type="checkbox"/> Venus <input type="checkbox"/> Mars <input type="checkbox"/> Jupiter <input type="checkbox"/> Saturn <input type="checkbox"/> Uranus <input type="checkbox"/> Neptune	
2.	<b>Drawing 1</b>	<b>Drawing 2</b>
3.		

## The Moon (32 points )

6 Points for pointing marked by the supervisor.



1. 2 points for drawing the moon  
2 points for drawing the lunar terminator properly.
2. 4 points for cardinal points
3. (2x3 = 6 points for maria, 2x3 = 6 points for craters)



4.

Age of the Moon (2 points)	8.3 days (8 - 8.5 days will gain full marks) (7.5 - 9 will gain half marks)
Hemisphere (Northern or Southern) from where the Moon is being watched (2 points)	Southern
Official IAU name of the constellation in which the Moon is located (2 points)	Taurus

The Planet (16 points)

1. Jupiter (2 points)
2. 4 points for drawing initial and final positions. 4 points for labelling them correctly.
3.  $1.5 \times 4 = 6$  for correct order.

Image 1 solution





Image 2 solution



Note: Students won't be writing the names. They will label them as A,B,C,D. The moon must have the same label in both images.

3.

Order:

**Ganymede**

**Callisto**

**IO**

**Europa**

